

## Literature References on Elements of Mixture Interpretation

### Mixture Principles & Recommendations

Buckleton, J.S., & Curran, J.M. (2008). A discussion of the merits of random man not excluded and likelihood ratios. *Forensic Science International: Genetics*, 2, 343-348.

Budowle, B., et al. (2009). Mixture interpretation: defining the relevant features for guidelines for the assessment of mixed DNA profiles in forensic casework. *Journal of Forensic Sciences*, 54, 810-821.

DNA Advisory Board (2000) Statistical and population genetic issues affecting the evaluation of the frequency of occurrence of DNA profiles calculated from pertinent population database(s) (approved 23 February 2000). *Forensic Science Communications*, July 2000. Available at: <http://www.fbi.gov/programs/lab/fsc/backissu/july2000/dnastat.htm>.

Gill, P., et al. (2006). DNA commission of the International Society of Forensic Genetics: Recommendations on the interpretation of mixtures. *Forensic Science International*, 160, 90-101. Available at <http://www.isfg.org/Publication;Gill2006>.

Gill, P., et al. (2008). National recommendations of the technical UK DNA working group on mixture interpretation for the NDNAD and for court going purposes. *Forensic Science International: Genetics*, 2, 76-82.

Morling, N., et al. (2007). Interpretation of DNA mixtures – European consensus on principles. *Forensic Science International: Genetics*, 1, 291-292.

Schneider, P.M., et al. (2006). Editorial on the recommendations of the DNA commission of the ISFG on the interpretation of mixtures. *Forensic Science International*, 160, 89-89.

Schneider, P.M., et al. (2009). The German Stain Commission: recommendations for the interpretation of mixed stains. *International Journal of Legal Medicine*, 123, 1-5. (originally published in German in 2006 -- *Rechtsmedizin* 16:401-404).

Stringer, P., et al. (2009). Interpretation of DNA mixtures—Australian and New Zealand consensus on principles. *Forensic Science International: Genetics*, 3, 144-145.

SWGDM (2010). SWGDAM interpretation guidelines for autosomal STR typing by forensic DNA testing laboratories. Available at [http://www.fbi.gov/hq/lab/html/codis\\_swgdam.pdf](http://www.fbi.gov/hq/lab/html/codis_swgdam.pdf).

Wickenheiser, R.A. (2006). General guidelines for categorization and interpretation of mixed STR DNA profiles. *Canadian Society of Forensic Science Journal*, 39, 179-216. Available at [http://raywickenheiser.com/pdf/wickenheiser\\_mixture\\_interpretation.pdf](http://raywickenheiser.com/pdf/wickenheiser_mixture_interpretation.pdf).

### Setting Thresholds

Currie, L. (1999). Detection and quantification limits: origin and historical overview. *Analytica Chimica Acta*, 391, 127–134.

Gilder, J.R., et al. (2007). Run-specific limits of detection and quantitation for STR-based DNA testing. *Journal of Forensic Sciences*, 52, 97-101.

Gill, P., et al. (2009). The *low-template-DNA* (stochastic) threshold -- its determination relative to risk analysis for national DNA databases. *Forensic Science International: Genetics*, 3, 104-111.

Gill, P. and Buckleton, J. (2010). A universal strategy to interpret DNA profiles that does not require a definition of *low-copy-number*. *Forensic Science International: Genetics*, 4, 221-227.

- Kaiser, H. (1970). Report for analytical chemists: part II. Quantitation in elemental analysis. *Analytical Chemistry*, 42, 26A-59A.
- Long, G.L., & Winefordner, J.D. (1983). Limit of detection: a closer look at the IUPAC definition. *Analytical Chemistry*, 55, 712A-724A.
- Miller J.C., & Miller J.N. (2005). Errors in instrumental analysis; regression and correlation in *Statistics for Analytical Chemistry*, Ellis Horwood and Prentice Hall, pp. 101-137.
- Mocak, J., Bond, A.M., Mitchell, S., & Scollary, G. (1997). A statistical overview of standard (IUPAC and ACS) and new procedures for determining the limits of detection and quantification: application to voltammetric and stripping techniques. *Pure and Applied Chemistry*, 69, 297-328.
- Rubinson, K.A., & Rubinson, J.F. (2000). Sample size and major, minor, trace, and ultratrace components. *Contemporary instrumental analysis*. Upper Saddle River: Prentice Hall, pp. 150–158.
- Tvedebrink, T., et al. (2009). Estimating the probability of allelic drop-out of STR alleles in forensic genetics. *Forensic Science International: Genetics*, 3, 222-226.

## Stutter Products

- Blackmore, V.L., et al. (2000). Preferential amplification and stutter observed in population database samples using the AmpFISTR Profiler multiplex system. *Canadian Society of Forensic Sciences Journal*, 33, 23-32.
- Gibb, A.J., et al. (2009). Characterisation of forward stutter in the AmpFISTR SGM Plus PCR. *Science & Justice*, 49, 24-31.
- Gill, P., et al. (1997). Development of guidelines to designate alleles using an STR multiplex system. *Forensic Science International*, 89, 185-197.
- Gill, P., et al. (1998). Interpretation of simple mixtures when artifacts such as stutters are present—with special reference to multiplex STRs used by the Forensic Science Service. *Forensic Science International*, 95, 213-224.
- Hill, C.R., et al. (2010). Concordance and population studies along with stutter and peak height ratio analysis for the PowerPlex((R)) ESX 17 and ESI 17 Systems. *Forensic Science International: Genetics*, (in press).
- Leclair, B., et al. (2004). Systematic analysis of stutter percentages and allele peak height and peak area ratios at heterozygous STR loci for forensic casework and database samples. *Journal of Forensic Sciences*, 49, 968-980.
- Mulero, J.J., et al. (2006). Characterization of the N+3 stutter product in the trinucleotide repeat locus DYS392. *Journal of Forensic Sciences*, 51, 1069-1073.
- Walsh, P.S., et al. (1996). Sequence analysis and characterization of stutter products at the tetranucleotide repeat locus vWA. *Nucleic Acids Research*, 24, 2807-2812.

## Stochastic Effects and Allele Dropout

- Balding, D.J., & Buckleton, J. (2009). Interpreting low template DNA profiles. *Forensic Science International: Genetics*, 4: 1-10.

Gill, P., et al. (2005). A graphical simulation model of the entire DNA process associated with the analysis of short tandem repeat loci. *Nucleic Acids Research*, 33, 632-643.

Puch-Solis, R., et al. (2009). Assigning weight of DNA evidence using a continuous model that takes into account stutter and dropout. *Forensic Science International: Genetics Supplement Series*, 2, 460-461.

Stenman, J., & Orpana, A. (2001). Accuracy in amplification. *Nature Biotechnology*, 19, 1011-1012.

Taberlet, P., et al. (1996). Reliable genotyping of samples with very low DNA quantities using PCR. *Nucleic Acids Research*, 24, 3189-3194.

Tvedebrink, T., et al. (2008). Amplification of DNA mixtures—missing data approach. *Forensic Science International: Genetics Supplement Series*, 1, 664-666.

Tvedebrink, T., et al. (2009). Estimating the probability of allelic drop-out of STR alleles in forensic genetics. *Forensic Science International: Genetics*, 3, 222-226.

Walsh, P.S., et al. (1992). Preferential PCR amplification of alleles: Mechanisms and solutions. *PCR Methods and Applications*, 1, 241-250.

## Peak Height Ratios

Bright, J.A., et al. (2010). Examination of the variability in mixed DNA profile parameters for the Identifiler multiplex. *Forensic Science International: Genetics*, 4, 111-114.

Buckleton, J. (2009). Validation issues around DNA typing of low level DNA. *Forensic Science International: Genetics*, 3, 255-260.

Buse, E.L., et al. (2003). Performance evaluation of two multiplexes used in fluorescent short tandem repeat DNA analysis. *Journal of Forensic Sciences*, 48, 348-357.

Debernardi, A., et al. (2010). One year variability of peak heights, heterozygous balance and inter-locus balance for the DNA positive control of AmpFISTR Identifiler STR kit. *Forensic Science International: Genetics*, (in press). DOI: 10.1016/j.fsigen.2010.01.020

Hill, C.R., et al. (2010). Concordance and population studies along with stutter and peak height ratio analysis for the PowerPlex® ESX 17 and ESI 17 Systems. *Forensic Science International: Genetics*, (in press).

Leclair, B., et al. (2004). Systematic analysis of stutter percentages and allele peak height and peak area ratios at heterozygous STR loci for forensic casework and database samples. *Journal of Forensic Sciences*, 49, 968-980.

Moretti, T.R., et al. (2001). Validation of short tandem repeats (STRs) for forensic usage: performance testing of fluorescent multiplex STR systems and analysis of authentic and simulated forensic samples. *Journal of Forensic Sciences*, 46, 647-660.

Moretti, T.R., et al. (2001). Validation of STR typing by capillary electrophoresis. *Journal of Forensic Sciences*, 46, 661-676.

Wallin, J.M., et al. (1998). TWGDAM validation of the AmpFISTR Blue PCR amplification kit for forensic casework analysis. *Journal of Forensic Sciences*, 43, 854-870.

## Estimating the Number of Contributors

- Brenner, C.H., et al. (1996). Likelihood ratios for mixed stains when the number of donors cannot be agreed. *International Journal of Legal Medicine* 109, 218-219.
- Buckleton, J.S., et al. (1998). Setting bounds for the likelihood ratio when multiple hypotheses are postulated. *Science & Justice* 38, 23-26.
- Buckleton, J.S., et al. (2007). Towards understanding the effect of uncertainty in the number of contributors to DNA stains. *Forensic Science International: Genetics*, 1, 20-28.
- Clayton, T.M., et al. (2004). A genetic basis for anomalous band patterns encountered during DNA STR profiling. *Journal of Forensic Sciences*, 49, 1207-1214.
- Egeland, T., et al. (2003). Estimating the number of contributors to a DNA profile. *International Journal of Legal Medicine*, 117, 271-275.
- Haned, H., et al. (2010). The predictive value of the maximum likelihood estimator of the number of contributors to a DNA mixture. *Forensic Science International: Genetics*, (in press).
- Haned, H., et al. (2010). Estimating the number of contributors to forensic DNA mixtures: does maximum likelihood perform better than maximum allele count? *Journal of Forensic Sciences*, (in press).
- Lauritzen, S.L., & Mortera, J. (2002). Bounding the number of contributors to mixed DNA stains. *Forensic Science International* 130, 125-126.
- Paoletti, D.R., et al. (2005). Empirical analysis of the STR profiles resulting from conceptual mixtures. *Journal of Forensic Sciences*, 50, 1361-1366.

## Mixture Ratios

- Clayton, T.M., et al. (1998). Analysis and interpretation of mixed forensic stains using DNA STR profiling. *Forensic Science International*, 91, 55-70.
- Cowell, R.G., et al. (2007). Identification and separation of DNA mixtures using peak area information. *Forensic Science International*, 166, 28-34.
- Evett, I.W., et al. (1998). Taking account of peak areas when interpreting mixed DNA profiles. *Journal of Forensic Sciences*, 43, 62-69.
- Frégeau, C.J., et al. (2003). AmpFISTR Profiler Plus short tandem repeat DNA analysis of casework samples, mixture samples, and nonhuman DNA samples amplified under reduced PCR volume conditions (25 microL). *Journal of Forensic Sciences*, 48, 1014-1034.
- Gill, P., et al. (1998). Interpreting simple STR mixtures using allelic peak areas. *Forensic Science International*, 91, 41-53.
- Perlin, M.W., & Szabady, B. (2001). Linear mixture analysis: a mathematical approach to resolving mixed DNA samples. *Journal of Forensic Sciences*, 46, 1372-1378.
- Wang, T., et al. (2006). Least-squares deconvolution: a framework for interpreting short tandem repeat mixtures. *Journal of Forensic Sciences*, 51, 1284-1297.

## Statistical Approaches

- Balding, D.J. (2005) *Weight-of-evidence for Forensic DNA Profiles*. John Wiley & Sons; see mixture section on pp. 101-110.
- Curran, J.M., et al. (1999). Interpreting DNA mixtures in structured populations. *Journal of Forensic Sciences*, 44, 987-995.
- Curran, J.M., & Buckleton, J. (2010). Inclusion probabilities and dropout. *Journal of Forensic Science*, 55, 1171-1173.
- Devlin, B. (1993). Forensic inference from genetic markers. *Statistical Methods in Medical Research*, 2, 241–262.
- Evett, I.W., et al. (1991). A guide to interpreting single locus profiles of DNA mixtures in forensic cases. *Journal of Forensic Science Society*, 31, 41-47.
- Evett, I.W., & Weir, B.S. (1998). *Interpreting DNA Evidence: Statistical Genetics for Forensic Scientists*. Sunderland, MA: Sinauer Associates.
- Fung, W.K., & Hu, Y.-Q. (2001). The evaluation of mixed stains from different ethnic origins: general result and common cases. *International Journal of Legal Medicine*, 115, 48-53.
- Fung, W.K., & Hu, Y.-Q. (2002). The statistical evaluation of DNA mixtures with contributors from different ethnic groups. *International Journal of Legal Medicine*, 116, 79-86.
- Fung, W.K., & Hu, Y.-Q. (2008). *Statistical DNA Forensics: Theory, Methods and Computation*. Wiley: Hoboken, NJ.
- Hu, Y.-Q., & Fung, W.K. (2003). Evaluating forensic DNA mixtures with contributors of different structured ethnic origins: a computer software. *International Journal of Legal Medicine*, 117, 248-249.
- Ladd, C., et al. (2001). Interpretation of complex forensic DNA mixtures. *Croatian Medical Journal*, 42, 244-246.
- Puch-Solis, R., et al. (2010). Calculating likelihood ratios for a mixed DNA profile when a contribution from a genetic relative of a suspect is proposed. *Science & Justice*, (*in press*).
- van Nieuwerburgh, F., et al. (2009). Impact of allelic dropout on evidential value of forensic DNA profiles using RMNE. *Bioinformatics* 25, 225-229.
- Van Nieuwerburgh, F., et al. (2009). RMNE probability of forensic DNA profiles with allelic drop-out. *Forensic Science International: Genetics Supplement Series*, 2, 462-463.
- Weir, B.S., et al. (1997). Interpreting DNA mixtures. *Journal of Forensic Sciences* 42, 213-222.

## Software

- Bill, M., et al. (2005). PENDULUM-a guideline-based approach to the interpretation of STR mixtures. *Forensic Science International*, 148, 181-189.
- Mortera, J., et al. (2003). Probabilistic expert system for DNA mixture profiling. *Theoretical and Population Biology*, 63, 191-205.

Oldroyd, N., & Shade, L.L. (2008) Expert assistant software enables forensic DNA analysts to confidently process more samples. *Forensic Magazine Dec 2008/Jan 2009*, 25-28; available at <http://www.forensicmag.com/articles.asp?pid=240>.

Perlin, M.W. (2006). Scientific validation of mixture interpretation methods. *Proceedings of Promega's Seventeenth International Symposium on Human Identification*. Available at <http://www.promega.com/geneticidproc/ussymp17proc/oralpresentations/Perlin.pdf>

## Probabilistic Genotyping Approach

Cowell, R.G., et al. (2010). Probabilistic expert systems for handling artifacts in complex DNA mixtures. *Forensic Science International: Genetics*, (in press).

Curran, J.M. (2008). A MCMC method for resolving two person mixtures. *Science & Justice*, 48, 168-177.

Perlin, M.W., & Sinelnikov, A. (2009). An information gap in DNA evidence interpretation. *PloS ONE*, 4(12), e8327.

Perlin, M.W., et al. (2009). Match likelihood ratio for uncertain genotypes. *Law, Probability and Risk*, 8, 289-302.

## General Information on Mixtures

Clayton, T., & Buckleton, J. (2005). Mixtures. Chapter 7 in *Forensic DNA Evidence Interpretation* (Eds.: Buckleton, J., Triggs, C.M., Walsh, S.J.), CRC Press, pp. 217-274.

Tomsey, C.S., et al. (2001). Case work guidelines and interpretation of short tandem repeat complex mixture analysis. *Croatian Medical Journal*, 42, 276-280.

Torres, Y., et al. (2003). DNA mixtures in forensic casework: a 4-year retrospective study. *Forensic Science International*, 134, 180-186.